

Nuclear Reactions for Astrophysics at NIF Using Radiochemistry “Thoughts”


Anna Hayes, P. Bradley, G.Hale, A. Heger, J. Jungman, R. Korzekwa
(Theory)

G. Grim, R. Rundberg, J. Wilhelmy
(Expt)

High Fluence of both Neutrons and Charged-Particles Induce Reactions off the line of stability

Allows multiple reactions on materials loaded in the shell
=> Reactions on short-lived unstable targets

- First neutron reactions to get off line of stability
(n,2n), (n,3n) or double (n,2n) for proton-rich
(n,γ) for neutron-rich
- Followed by charged-particle reactions of interest
e.g., (p,γ), (p,α), ...



Caveat: NIF design physicists warn that doping is possible *iff* achieve robust ignition

Currently Developing Diagnostic for Mix and Instabilities in Ignited NIF Capsules


Scheme based on using double reactions:

$$\begin{array}{lcl} n+t & \longrightarrow & t^*(\text{knock-on}) \\ t+\text{shell} & \longrightarrow & X(\beta^-) \end{array}$$

Detect



Triton interactions with shell material ~100mb. Mix diagnosed by observing beta decays of resulting isotopes, after target disassembly.

Triton + ^9Be ablator (t, α) reactions distinguish different types of mix.



NIF full Yield - ${}^8\text{Li}$ production

No Mix	Chunk Mix	Atomic Mix
1×10^{12}	3×10^{12}	1×10^{14}



Debris collector
& β counter.

Thoughts on using similar techniques for Nuclear Astrophysics

- How far off the line of stability can we go?
- What can we do that we can't do at a radioactive ion beam facility?
- How do we detect the reactants?
- How many reactions do we need for collection/detection?
- Which reactions are feasible?
- How do extrapolate from NIF flux-average the cross section to those needed of astrophysicist?

Physics Parameters for Ignited Capsule

Examined a high yield DT capsule with ${}^9\text{Be}(.9\%\text{Cu})$ ablator/shell

Clean yield 16.65 MJ \Rightarrow 6×10^{18} DT reactions, $\sim 6 \times 10^{18}$ neutrons

$$\langle \rho r \rangle_{\text{shell}} = 0.25 \text{ g/cm}^2$$
$$\text{range} = .1 - .6 \text{ g/cm}^2$$


$$\langle \rho r \rangle_{\text{capsule}} = 1.2 \text{ g/cm}^2$$

Also examined mixing between shell and DT
Mix varied as a parameter - the mixing length

$$L = \alpha \int |u_{DT}(r, t) - u_{\text{shell}}(r, t)| dt$$

$$\alpha = 0.06 \Rightarrow Y = 9.5 \text{ MJ}$$

$$\alpha = 0.12 \Rightarrow Y = 5.2 \text{ MJ}$$




Reaction estimates scale with yield and $\langle \rho r \rangle$

Neutron Spectrum


Neutron spectra at DT fuel/shell interface

Different yields and shell/fuel mixing result in similarly shaped spectra

10% < 14 MeV peak
0.3% > 14 MeV peak (RIFs)



Neutron Reactions



(n,2n) reactions best potential
Typical cross sections $\sim 1\text{b}$ at 14 MeV

Single (n,2n) $\sim 10^{16}$ reactions
Double (n,2n) $\sim 10^{12} - 10^{13}$ reactions

Some heavy nuclei (n,3n) $\sim 1\text{b}@ 14\text{ MeV}$

However, for many nuclei of interest
neutron binding is high
 \Rightarrow (n,2n) threshold $> 14\text{ MeV}$

e.g., ^{32}S ; ^{24}Mg , ^{16}O ; ^{19}F ,

$10^{16} \longrightarrow \sim 10^{13}$ or less



Other Neutron Reactions also Possible

➤ The so-called direct-semi-direct contribution to **neutron capture** yields a capture cross sections $\sim 1\text{mb}$ @ 14 MeV
 $\Rightarrow \sim 5 \times 10^{13}$ reactions

➤ **Inelastic (n,n')** scattering to isomers
 $\Rightarrow 3 \times 10^{15}$


➤ **Threshold for (n,3n) is often too high to get** a significant number of reactions

➤ Double (n,2n) often better



Charged Particle Fluences

Main Reactions




Produce large fluence of energetic (MeV)
p, d, t, ${}^3\text{He}$, ${}^4\text{He}$


Can retain significant fraction of this energy
upon reaching the ablator

Thermal Broadening


$d+t \rightarrow n+{}^4\text{He}$



$d+d \rightarrow p+{}^3\text{H}$



Charged-Particle Fluence in the Shell



Energetic p and d can traverse the shell

At the fuel-shell interface


n	6×10^{18}	14.0 MeV
α	3×10^{18}	0.4 MeV
p	1×10^{17}	2.3 MeV
t	4×10^{17}	0.5 MeV
d	3×10^{17}	1.2 MeV

Outer surface of ablator

n	5×10^{18}	14.0 MeV
α	~ 0	0
p	3×10^{15}	~ 50 keV
t	few	~ 10 keV
d	1×10^{16}	~ 50 keV

Example

Want to measure (p,γ) and (p,α) on unstable waiting point nucleus



Non-thermal charged particle Reactions

- A significant fraction of charged-particle reactions can take place before thermalizing
- Energetic (MeV) cross-sections can be orders of magnitude larger than thermal
- Stopping and Non-Maxwellian component of charged-particle spectrum need to be understood in detail
- Probably need full Monte Carlo treatment of charged-particle transport & reaction

Number of charged-particle Reactions Produced depends on energy & stopping

Stopping Length:

$$L \simeq \frac{2\theta_e^{3/2} E_0^{1/2}}{C_R n_e}$$

Probability of particle with energy E reacting:

$$P(E) \approx \hat{n} \int_0^E dE' \frac{\sigma(E)}{-(dE/dx)_{shell}} \approx 2 \frac{n\theta_e^{3/2}}{n_e C} \int dE' \frac{\sigma(E)}{\sqrt{E}}$$

Need to integrate this over
energy distribution of charged-particles
in the shell

Ratio of density of unstable targets
to electrons in the shell $\sim 10^{-5} - 10^{-6}$

Count Estimates

Dope shell with A (stable) (say 33%)

- $A(n,2n)A-11(n,2n)A-2$
=> $\sim 10^{12}$ produced, if both (n,2n) cross section 1b


10^{17} protons at fuel/shell interface $\langle E_p \rangle \sim 2.3$ MeV

3×10^{15} protons make it to outer surface of shell $\langle E_p \rangle \sim 50$ keV

- **Get $\sim 10^5$ reactions assuming a 10mb cross section**
- *(integration over p spectrum and stopping preliminary)*
- Simplified estimate: $N_p \langle \rho r \rangle (A-2/A \text{ atoms}) \sigma = 5 \times 10^4$ reactions


Up Scattered Charged-Particles Observed at Omega

DT Neutron



Stopping Length

$$\frac{d\psi_t}{dE} = \frac{\theta_e^{\frac{3}{2}} \psi_n n_t}{c_R n_e \sqrt{E}} \sigma_{nt}(\geq E)$$



Produce p,d,t with $E > 10$ MeV


Escape the capsule and detected

Emitted charged-particles are Strong constraint on predicted charged-particle transport, stopping, etc.

$$\langle \rho r \rangle_{\text{shell}} \sim 35 \text{ mg/cm}^2$$

Knock-on tritons at Omega predicted to loose about 3.45 MeV traversing the shell


In reasonable agreement with experiment



Ongoing Experimental Program Beta-Mix

MeV Triton interaction with shell material ~100mb.

Observe beta decays of resulting isotopes, after target disassembly.




Ignited d-t
Fuel

Main Reactions

NIF: ${}^9\text{Be}(t, \alpha){}^8\text{Li}$ (β^- , 840ms)
 ${}^9\text{Be}(t, p){}^{11}\text{Be}$ (β^- , 13.8s)

Omega: ${}^{13}\text{C}(t, p){}^{15}\text{C}$ (β^- , 2.5s)
 ${}^{13}\text{C}(t, \alpha){}^{12}\text{B}$ (β^- , 21 ms)


All high-energy β 's ~ 10 MeV



Debris collector
& β counter.

DEBRIS COLLECTION CHALLENGES

- Need high efficiency and sufficient solid angle.
- Induced radioactivity must be low.
- Must survive the radiation and shock wave.
- Must be able to assay debris fraction i.e., bomb fraction.
- Overcome hohlraum issues



430 ns
2900 ns


160 ns
1000 ns

0.5 ns
3.5 ns

Omega (15 cm)
NIF (1 m?)

Progress using the Omega Laser Facility


- Fielded β -detector with 30 ms recovery
- Observed β -decays with half-lives as short as ^{12}B



- Preliminary measurement of 300 μb , $^{12}\text{C}(n,p)^{12}\text{B}$ cross section at 14 MeV.
- Observed $^{15}\text{C}(2.45 \text{ sec } \beta^-)$ from the $^{15}\text{N}(n,p)$ reaction - extracting cross section
- First debris collection experiments fielded using labeled ^{13}C targets.
- Debris collector mass measurement sensitivity better than 0.5 nanograms.

Reactions in the Hohlraum

- Baseline design is cylindrical
75%U, 25%Au cocktail
- Wall thickness is $30\mu\text{m}$, of which $7\mu\text{m}$ is the cocktail
- Possible to use other materials for the wall backing without affecting the yield



10^{19} neutrons \Rightarrow $\sim 10^{15}$ reactions for 1b cross section
 $\sim 10^{11}$ for double reactions

Summary

- Fluence of neutrons and charged particles suggest that double and triple reactions produce significant yields
- Reactions of interest very challenging - triple reactions maybe too small
- Significant progress being made to develop experimental techniques at Omega
- Can measure reaction products with lifetimes ~ tens msec
- Energetic of charged-particles traversing the shell high
- Need accurate tools to extract the desired thermal cross sections from flux-averaged measurements
- Reactions in the hohlraum may be a possibility